

IL-12/NKSF/CLMF/TSF, Human, Recombinant

货号：PCK302

产品信息

别名	Interleukin-12, NKSF, Cytotoxic Lymphocyte Maturation Factor (CLMF), TSF
物种	Human
表达宿主	HEK293
序列信息	IWELKKDVYVVELDWYPDAPGEMVVLTCDTPEEDGITWTL DQSSEVLGSG KTLTIQVKEFGDAGQYTCHKGGEVLSHSLLLHKKEDGIWSTDILKDQKEP KNKTFLRCEAKNYSGRFTCWWLTTISTDLTFSVKSSRGSSDPQGVTCGAAT LSAERVRGDNKEYEYSVECQEDSACPAAEESLPIEVMVDAVHKLKYENYTS SFFIRDIKPDPPKNLQLKPLKNSRQVEVSWEYPDTWSTPHSYFSLT
检索号	P29459(p35), P29460(p40)
分子量	59.55 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to induce IFN gamma secretion in PHA-activated human peripheral blood lymphocytes (PBMC). The ED50 for this effect is 0.05-0.2 ng/mL.

产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Active IL-12 is a p70 disulfide-linked dimer composed of p35 and p40 subunits. The protein is a pleiotropic cytokine produced primarily by antigen presenting cells and has multiple effects on T lymphocytes and natural killer cells in terms of stimulating cytotoxicity, proliferation, production of other cytokines and Th1 subset differentiation.

SDS-PAGE

