

LIF, Mouse, Recombinant

货号 : PCK307

产品信息

别名	leukemia inhibitory factor, Differentiation-stimulating factor, D factor, Melanoma-derived LPL inhibitor.
物种	Mouse
表达宿主	E.coli
序列信息	SPLPITPVNATCAIRHPCHGNLMNQIKNQLAQLNGSANALFISYYTAQGEPF PNNVEKLCAPNMTDFPSFHGNGTEKTKLVELYRMVAYLSASLTNITRDQK VLNPTAVSLQVKLNATIDVMRGLLSNVLCRLCNKYRVGHVDVPPVPDHSD KEAFQRKKLGCQLLGTYKQVISVVVQAF with polyhistidine tag at the N-terminus.
检索号	P09056.1
分子量	20.68 kDa
标签	His-tag at the N-terminus
生物活性	Measure by its ability to induce IL-6 secretion in M1 cells. The ED50 for this effect is <0.5 ng/mL. The specific activity of recombinant mouse LIF is > 2 x 10 ⁶ IU/mg.

产品特性

纯度	>95% as determined by SDS-PAGE. Ni-NTA chromatography.
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

LIF, a pleiotrophic factor, is identified in multiple cell types, including T cells, myelomonocytic lineages, fibroblasts, liver, heart and melanoma. LIF is capable of promoting long-term maintenance of embryonic stem cells by inhibiting spontaneous differentiation. In addition, LIF also have abilities including stimulation of differentiation of cholinergic nerves, the stimulation of acute phase protein synthesis by hepatocytes, and suppression of adipogenesis by supressing the lipoprotein lipase in adipocytes.

SDS-PAGE

