

FGF-9/ HBGF-9/ GAF, Human, Recombinant

货号 : PCK138

产品信息

别名	Fibroblast Growth Factor 9; FGF-9; Glia-Activating Factor; GAF; Heparin-Binding Growth Factor 9; HBGF-9; FGF9
物种	Human
表达宿主	E.coli
序列信息	Met1-Ser208
检索号	P31371
分子量	23.44 kDa
生物活性	Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED50 for this effect is 1-5 ng/ml.

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 220mM Sucrose , 0.02% Tween 80 , pH 6.0.




复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


背景介绍

Fibroblast Growth Factor 9 (FGF-9) belongs to the Fibroblast Growth Factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF-9 plays an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration. In addition, FGF-9 may have a role in glial cell growth and differentiation during development, gliosis during repair and regeneration of brain tissue after damage, differentiation and survival of neuronal cells, and growth stimulation of glial tumors.

SDS-PAGE



生物活性



Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED50 for this effect is 1-5 ng/ml.

