

IL-10/ IL10A, Human, Recombinant

货号 : PCK169

产品信息

别名	Interleukin-10; IL-10; Cytokine synthesis inhibitory factor; CSIF; IL10; RP11-262N9.1; IL10A; MGC126450; MGC126451; TGIF
物种	Human
表达宿主	Human Cells
序列信息	Ser19-Asn178
检索号	P22301
分子量	18.6 kDa
生物活性	Loaded Human IL-10RA-His on HIS1K Biosensor, can bind Human IL-10 with an affinity constant of 1 pM as determined in BLI assay.

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.




复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


背景介绍

Interleukin 10 (IL10), also known as Cytokine synthesis inhibitory factor (CSIF), is a secreted Protein and belongs to the IL-10 family. IL-10 is secreted by many activated hematopoietic cell types as well as hepatic stellate cells, keratinocytes, and placental cytotrophoblasts. IL-10 is an anti-inflammatory TH2 Cytokine that has a critical role in limiting the immune response to pathogens to prevent host damage. As IL-10 is produced in several T helper populations, it is proposed that it provides a feedback loop to limit the effector functions of macrophages and DCs on T cells. Once expressed, IL-10 signals through the IL-10 Receptor (IL-10R) to activate STAT3. As IL-10 is a strong inhibitor of inflammation, it has become a viable biomarker for various diseases and conditions as well as a therapeutic molecule for certain conditions. In addition to elevated levels in parasitic infection, high expression levels of IL-10 are also found in retroviral infections inducing immunodeficiency. The immunosuppressive properties of IL-10 suggest a possible clinical use of IL-10 in suppressing rejections of grafts after organ transplantations.

SDS-PAGE



生物活性



Loaded Human IL-10RA-His on HIS1K Biosensor, can bind Human IL-10 with an affinity constant of 1 pM as determined in BLI assay.

