

IL-10/B-TCGF/CSIF/TGIF, Mouse, Recombinant

货号 : PCK248

产品信息

别名	B-TCGF, CSIF, TGIF
物种	Mouse
表达宿主	E.coli
序列信息	MSRGQYSREDNNCTHFPVGQSHMLLELRTAFSQVKTFQTKDQLDNILLTD SLMQDFKGYLGCQALSEMIQFYLVEMPQAEKHGPEIKEHLNSLGEKLKTL RMRLRRCHRFLPCENKSKAVEQVKSDFNKLQDQGVYKAMNEFDIFINCIEA YMMIKMKS with polyhistidine tag at the C-terminus.
检索号	P18893.2
分子量	19.69 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to induce MC/9-2 cells proliferation. The ED50 for this effect is <1 ng/mL. The specific activity of recombinant mouse IL-10 is > 1 x 10 ⁶ IU/mg.

产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Interleukin 10 (IL-10), also known as human cytokine synthesis inhibitory factor (CSIF), is an anti-inflammatory cytokine. In humans, interleukin 10 is encoded by the IL10 gene. IL-10 signals through a receptor complex consisting of two IL-10 receptor-1 and two IL-10 receptor-2 proteins. Consequently, the functional receptor consists of four IL-10 receptor molecules. IL-10 binding induces STAT3 signalling via the phosphorylation of the cytoplasmic tails of IL-10 receptor 1 + IL-10 receptor 2 by JAK1 and Tyk2 respectively.

SDS-PAGE

