

IL-12 p40/IL-12B/CLMF p40, Mouse, Recombinant

货号: PCK249

产品信息

别名	Interleukin-12 subunit beta, IL-12 subunit p40, IL-12B, Cytotoxic Lymphocyte Maturation Factor 40 kDa subunit (CLMF p40), NK cell Stimulating Factor Chain 2
物种	Mouse
表达宿主	E.coli
序列信息	MWELEKDVYVVEVDWTPDAPGETVNLTCDTPEEDDITWTSDQRHGVIGSG KTLTITVKEFLDAGQYTCHKGGETLSHSHLLHKKENGIWSTEILKNFKNKT FLKCEAPNYSGRFTCSWL VQRNMDLKFNKSSSSSPDSRAVTCGMASLSAE KVTLDQRDYEKYSVSCQEDVTCPTAEETLPIELALEARQQNKYENYSTSFFI RDIKPDPPKNLQMKPLKNSQVEVSWEYPDSWSTPHSYFSLKFFV
检索号	P43432.1
分子量	36.60 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to induce proliferation in T-cell enriched PBMC. The ED50 for this effect is <0.3 ng/mL.

产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Active IL-12 is a p70 disulfide-linked dimer composed of p35 and p40 subunits. The protein is a pleiotropic cytokine produced primarily by antigen presenting cells and has multiple effects on T lymphocytes and natural killer cells in terms of stimulating cytotoxicity, proliferation, production of other cytokines and Th1 subset differentiation.

SDS-PAGE

