

IL-23/ NKSF2, Mouse, Recombinant

货号 : PCK256

产品信息

别名	SGRF; IL-23p19; CLMF p40; IL-12 subunit p40; NKSF2
物种	Mouse
表达宿主	Human Cells
序列信息	Val22-Ala196&Met23-Ser335
检索号	Q9EQ14&P43432
分子量	19.7&5.8 kDa

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of PBS, 5% Trehalose ,5% Mannitol, 0.01% Tween80, pH7.4.



复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

Interleukin 23 (IL-23) is a heterodimeric Cytokine composed of two disulfide-linked subunits, a p19 subunit that is unique to IL-23, and a p40 subunit that is shared with IL-12. The p19 subunit has homology to the p35 subunit of IL-12, as well as to other single chain Cytokines such as IL-6 and IL-11. The p40 subunit is homologous to the extracellular domains of the hematopoietic Cytokine Receptors. Although p19 is expressed by activated macrophages, dendritic cells, T cells, and endothelial cells, only activated macrophages and dendritic cells express p40 concurrently to produce IL-23. IL-23 has biological activities that are similar to, but distinct from IL-12. Both IL-12 and IL-23 induce proliferation and IFN-gamma production by human T cells. While IL-12 acts on both naive and memory human T cells, the effects of IL-23 is restricted to memory T cells.

SDS-PAGE

